

COVID Yellow: Moderate Risk of COVID-19 Spread

- There is a downward trajectory in numbers of COVID positive cases OR percent positive cases due to community or regional spread
- There is a downward trajectory in levels of Flu-Like Symptoms OR COVID like syndromic cases in Buffalo, Dawson, Franklin, Gosper, Harlan, Kearney, and Phelps Counties or regionally
- There is limited community spread within Buffalo, Dawson, Franklin, Gosper, Harlan, Kearney, and Phelps Counties or regionally
- COVID has a small impact on the local healthcare system capacity
- PPE supplies are adequate for critical healthcare workers and community needs
- Rapid COVID testing is available for all suspected cases and focused asymptomatic surveillance.
- TRPHD is able to conduct investigation and contact tracing for new cases and associated contacts without assistance
- Critical medical equipment is readily available

General Public	Outside the Home – Work, Social, In Public	At Home
Physical Distancing	<ul style="list-style-type: none"> • Consider staying at home most of the time, with caution for non-essential travel and work • Distance at least 6 feet from anyone outside the home • Work from home if possible • Individual outdoor activities such as walking, biking, etc. are acceptable • Cautious expansion of interactions with others, outdoor activities preferred • Gatherings only with modifications for COVID-19 • Events only as guided by TRPHD 	<ul style="list-style-type: none"> • No distancing for people without symptoms, unless they are under quarantine as a result of being exposed to someone who is infected • Self-isolation of symptomatic persons • Outdoor activities with distancing are acceptable.
Face Covering	<ul style="list-style-type: none"> • Face coverings suggested when unable to distance • Face coverings for symptomatic people 	<ul style="list-style-type: none"> • No face coverings for people without symptoms, unless they are under quarantine as a result of being exposed to someone who is infected • Face coverings for symptomatic people
Hand Washing	<ul style="list-style-type: none"> • Frequently wash hands for at least 20 seconds, especially after touching high contact surfaces, sneezing/coughing/touching face, or before eating • Use hand sanitizer when handwashing is inaccessible or infeasible 	<ul style="list-style-type: none"> • Frequently wash hands for at least 20 seconds, especially after touching high contact surfaces, sneezing/coughing/touching face, or before eating
Illness Monitoring	<ul style="list-style-type: none"> • If ill with Flu-like or COVID-like symptoms, Stay at Home • Minimize contact with symptomatic people • Daily temperature checks at work 	<ul style="list-style-type: none"> • Monitor for COVID-like symptoms
Disinfecting	<ul style="list-style-type: none"> • Avoid bare hand contact with any high touch surface • Use barrier such as paper towel or clothing when bare hand contact is unavoidable • Wash hands or apply sanitizer after touching high contact surfaces • Enhanced disinfection, especially for high contact surfaces (door/other handles, light switches, bathroom fixtures) 	<ul style="list-style-type: none"> • Enhanced disinfection, especially for high contact surfaces (door/other handles, light switches, bathroom fixtures)
At-Risk or Vulnerable Population	<p>For those over age 65, people with underlying health conditions, and other populations at heightened risk from COVID-19</p> <ul style="list-style-type: none"> • Stay home • Rely on help for needs outside the home (groceries, medications, etc.) • Distance from those working outside of the home 	